



Acute toxicity in two edible marine gastropods with different sensitivity to Tributyltin

MÓNICA ANGELINA PRIMOST^{1*}, SEBASTIÁN GIULIANELLI¹, MÓNICA NOEMI GIL^{2,3} & GREGORIO BIGATTI^{1,3}

¹LARBIM - Centro Nacional Patagónico – CONICET. Bvd. Brown 2915 (U9120ACV), Puerto Madryn, Chubut. Argentina.

²LOQyCA - Centro Nacional Patagónico – CONICET.

³Universidad Nacional de la Patagonia San Juan Bosco (UNPSJB), Facultad de Ciencias Naturales, Bvd. Brown 3051, Puerto Madryn, Chubut, Argentina.

*Corresponding author: primost@cenpat-conicet.gob.ar

Phone: +54 280 488-3184 int 1343 and Fax: +54 0280 4883543

Abstract

Buccinanops globulosus and *Trophon geversianus* are high and low sensitive to TBT. LD50 was 5.41 and 6.21 µgTBT.g⁻¹bw for *B. globulosus* and *T. geversianus* respectively, denoting high resistance to TBT acute exposure. These results can be used for future imposex induction experiments.

Keywords: mollusks, toxicity assays, TBT, environmental risk, imposex incidence, Patagonia.

Resumen: Toxicidad aguda del Tributilestano (TBT) en dos gasterópodos marinos comestibles. *Buccinanops globulosus* y *Trophon geversianus* son indicadores de alta y baja sensibilidad al TBT. La DL₅₀ fue de 5,41 y 6,21 µg TBT.g⁻¹ph para *B. globulosus* y *T. geversianus* respectivamente, denotando elevada resistencia a la exposición aguda de TBT. Estos resultados pueden ser útiles en futuros experimentos de inducción.

Palabras clave: moluscos, ensayos de toxicidad, TBT, riesgo ambiental, imposex, Patagonia.

Biological effects of chemicals and measurements of toxicity in impacted environments can be assessed by bioassays (Lee *et al.* 2005). Tributyltin (TBT) is a highly toxic compound to organisms (Thain *et al.* 1987, de Mora & Pelletier 1997, Evans 1999, Meador 2000, among others) and the analysis of this contaminant has been focused on the chronic effects caused by long time of exposure to polluted environments (Moore *et al.* 1991, Foale 1993, Ruiz *et al.* 1994, Pessoa *et al.* 2001, Antizar-Ladislao 2008). In marine invertebrates, particularly mollusks, sublethal effects such as imposex development (Gibbs & Bryan 1986) or

shell thickening in gastropods and bivalves for commercial purpose (Ruiz *et al.* 1995, Horiguchi *et al.* 1998, Bandow & Weltje 2012, among others) have been reported as consequence of TBT exposure.

The caenogastropod *Buccinanops globulosus* (Kiener, 1834) belonging to the family Nassariidae is an endemic species of the South Western Atlantic Ocean inhabiting sandy or muddy bottoms of shallow waters (Pastorino 1993). The genus has been affected by imposex phenomenon in all harbors and high maritime traffic areas from Argentina (Penchaszadeh *et al.* 2001, Bigatti *et al.* 2009,

Averbuj & Penchaszadeh 2010). On the other hand, the species *Trophon geversianus* (Pallas, 1774) belonging to Muricidae is widely distributed along the Atlantic and Pacific coasts of South America (Castellanos & Landoni 1993, Griffin & Pastorino 2005, Pastorino 2005), inhabiting hard bottoms. Muricids are the most used caenogastropods worldwide in monitoring studies due its high sensitivity to TBT (Gibbs & Bryan 1987, Axiak *et al.* 1995, Oehlmann *et al.* 1996). However, imposex development is very low in this species along the Argentinean coasts, even in harbor areas with TBT contamination (Bigatti *et al.* 2009). In contrast, *B. globulosus* has been classified as a good indicator of TBT contamination due to its high imposex levels even at low environmental concentrations (Bigatti *et al.* 2009). This edible species is also an expanding resource for artisanal fisheries in northern Patagonia, where it is locally consumed (Narvarte 2006, Averbuj *et al.* 2014) while *T. geversianus* is not consumed yet in the region (Bigatti *et al.* in press). However in Chile, *T. geversianus* is well consumed and commercialized (Leiva & Castilla 2002, González *et al.* 2007). Tributyltin concentrations up to 174.81 ngSn.g⁻¹dw were recently measured in coastal sediments of Nuevo gulf, Patagonia (Bigatti *et al.* 2009, del Brío [*unpublished data*]). At the same time, Bigatti *et al.* (2009) and then Primost (*unpublished data*) found incidence of imposex and negative reproductive effects on the affected populations. Despite of the TBT worldwide prohibition since year 2008 (IMO 2002), the contaminant is even found in marine environments (Antizar-Ladislao 2008, Zhang *et al.* 2013) and the populations of marine gastropods and other species could be at risk of physiological failures or local extermination.

To determine acute toxicity by median lethal dose (LD₅₀ 96h) assay, we injected both *Buccinanops globulosus* and *Trophon geversianus* with different doses of TBT. These results will be used as reference doses in future imposex induction experiments. Seawater and marine gastropods were collected from Cerro Avanzado beach (42°49'37.66"S; 64°51'29.19"W), a zone with scarce maritime traffic and very low imposex incidence for *B. globulosus* and *T. geversianus* (Bigatti *et al.* 2009, Primost [*unpublished data*]). Tests were performed with mature specimens (sizes over 18 mm for males and 30 mm for females [*unpublished data*]). They were identified and classified by observing the ventral pedal gland in females crawling over glass aquaria. Glass aquaria (20x30x20 cm) with 18 liters

of filtered seawater, also collected from Cerro Avanzado beach, were used in the tests. Water temperature (Collin & Spangler 2012) and salinity (35 mg/L) were controlled during the experiments using an OAKTON 600 series waterproof portable meter kit. A photoperiod of 12 h light/dark was regulated with a timer. Internal circulation inside each aquarium and constant aeration were provided to ensure organisms survival in laboratory conditions, but without recirculation among aquaria. Gastropods were acclimated for one week in each aquarium and starved before and during the experiment. Total body weight (0.1 g precision) of each gastropod was determined before injection in the laboratory. Exposure to TBT was performed by injecting liquid doses in the foot with a Hamilton micro syringe of 10 µl. Doses to be injected (µg of TBTCI per gram of wet body weight) were selected following previously published reports (Oberdörster & McClellan-Green 2000, Oberdörster & McClellan-Green 2002, Santos *et al.* 2005, Horiguchi *et al.* 2007, Garaventa *et al.* 2008) as well as own preliminary assays (0.1 and 0.5 µgTBTCI.g⁻¹ bw) using the same species (*unpublished data*). They were prepared from a stock solution of TBT chloride (Sigma Aldrich®) in absolute ethanol (960 µg of TBTCI per µl⁻¹). The relationship between total weight and wet body mass weight (bw) (Primost *unpublished data*) was used to calculate the injection volume of TBT per organism and are shown as supplementary data. For the test, eight groups of ten individuals each were conformed. Six treatments (1; 3; 5; 8; 10 and 20 µgTBTCI.g⁻¹bw) and two controls (without injection and with injection of 7 µl of absolute ethanol) were performed for both gastropod species. Size and weight parameters for gastropods in each treatment are shown in Table 1.

Once injected, gastropods were daily monitored. Total number of dead gastropods was recorded 96 hours later in each treatment. Lethal median dose LD₅₀ 96h and confidence intervals (95%) were calculated by XLStat 2012®-XLSTAT-Dose software for each species. Probit method (Finney 1971) with 100 iterations and 0,000001 of convergence was used.

Gastropods mortality was not observed in the two settled controls during the experiment for each species. No mortality was registered in *Trophon geversianus* in the dose of 1 µg while 10 % mortality occurred for the 3 µg TBTCI.g⁻¹bw dose. For doses higher than 5 µg TBTCI.g⁻¹bw, the proportion of dead gastropod increased for both species. Mortality was increased to 100% in the higher dose (20 µg

TBT.g⁻¹bw) for both *B. globulosus* and *T. geversianus* (Fig.1). Imposex development at the end

of the experiment (96 h) was analyzed and didn't find imposex induction in all treatments (Table 1).




Figure 1. Dose-response curves for *Trophon geversianus* and *Buccinanops globulosus*.

The lethal median dose calculated for 96h was 5.41 µg TBT.g⁻¹bw with 3.37 and 8.60 µg TBT.g⁻¹b as 95% confidence intervals in *B. globulosus*, and 6.21 µg TBT.g⁻¹bw with 4.61 and 8.08 µg TBT.g⁻¹bw as 95% confidence intervals in *T. geversianus*.

Reports of TBT lethal doses in marine gastropods and bivalves are scarce although acute effects using water dilutions of TBT have been reported (Morris 1987, Thain *et al.* 1987, Waldock *et al.* 1987, Bushong *et al.* 1988). The results obtained in our study with adult specimens indicated that TBT presented low acute toxicity for *Buccinanops globulosus* and *Trophon geversianus* compared with doses used to imposex induction (sublethal effects) in other species (Oberdörster & McClellan-Green 2000, Santos *et al.* 2005). Only for the highest dose (20 µgTBT.g⁻¹bw), 100% of mortality was recorded. Despite *T. geversianus* showed a higher value of LD₅₀ than *B. globulosus*, the overlapped confidence intervals do not allow ensuring a difference that can explain the different sensitivity to imposex development between muricids and nassarid (Bigatti *et al.* 2009). However, at low dose of 1 µg TBT.g⁻¹bw, also used by others authors in imposex induction experiments (Castro *et al.* 2007, Horiguchi *et al.* 2008), *B. globulosus* presented 20% of mortality while no *T. geversianus* died. Probably the differences in imposex incidence could be triggered by low concentrations of TBT in *B. globulosus*, and 1 µg TBT.g⁻¹bw could be the best starting dose in future imposex induction experiments in *T. geversianus*. Studies of imposex induction in caenogastropods have been conducted by injecting TBT into the foot region (Oberdörster &

McClellan-Green 2002, Santos *et al.* 2005) and even, it has been proposed that the pedal ganglia contains a penis morphogenic factor, which can interact directly to induce imposex development (Oberdörster & McClellan-Green 2000, Oberdörster *et al.* 2005). In this study, TBT was injected into the foot region in the same way that induction studies in other species of gastropods (Spooner *et al.* 1991, Oehlmann *et al.* 2000, McClellan-Green *et al.* 2006) to avoid the effect of adsorption of TBT on the walls of the aquarium (Laughlin *et al.* 1996, Panagoula *et al.* 2002). We assume that the whole TBT dose entered the muscular tissues of the gastropod. As the present study only focused on the short term (acute) effects of TBT, chronic and sublethal doses should be investigated on both *B. globulosus* and *T. geversianus* to better understand the chronic impacts and imposex development mediated by TBT. Acute toxicity tests conducted under controlled laboratory conditions are effective for exploring the worst situation of exposure (Matthiessen 2013) but provides no information about what actually occurs in the environment. In particular, interactions such as synergism or antagonism with other contaminants, as trace elements and polyaromatic hydrocarbons reported in Nuevo gulf (Gil *et al.* 2006, Commendatore & Esteves 2007, Giarratano *et al.* 2013, Giarratano *et al.* 2014, Gil *et al.* 2015, Primost *unpublished data*) may increase or reduce the tested effect. In this sense, experiments of imposex induction by TBT will be necessary in order to certainly corroborate the minimum dose to induce imposex in the species *B. globulosus* and *T. geversianus*.

Table 1. Size, weight (mean \pm standard deviation) and imposex parameters in *B. globulosus* and *T. geversianus* in each treatment

Treatment	Total size (mm)	Total weight (g)	Body mass weight (g)	Imposex % after 96 h
<i>Buccinanops globulosus</i>				
Without injection	36.99 \pm 2.00	9.23 \pm 1.65	6.32 \pm 0.72	0
7 μ l ethanol	41.92 \pm 0.83	12.04 \pm 0.74	5.64 \pm 0.36	0
1 μ g TBT.g ⁻¹	38.72 \pm 2.09	10.59 \pm 1.53	4.93 \pm 0.75	0
3 μ g TBT.g ⁻¹	40.39 \pm 1.28	11.99 \pm 0.98	4.05 \pm 0.48	0
5 μ g TBT.g ⁻¹	42.05 \pm 1.30	12.47 \pm 0.96	4.11 \pm 0.47	0
8 μ g TBT.g ⁻¹	39.14 \pm 1.73	10.79 \pm 1.15	5.02 \pm 0.57	0
10 μ g TBT.g ⁻¹	39.91 \pm 2.25	11.49 \pm 1.65	7.13 \pm 0.81	0
20 μ g TBT.g ⁻¹	39.27 \pm 2.33	10.99 \pm 1.60	7.35 \pm 0.79	0
<i>Trophon geversianus</i>				
Without injection	27.37 \pm 0.94	3.62 \pm 1.12	1.18 \pm 0.12	0
7 μ l ethanol	27.95 \pm 0.90	3.85 \pm 1.06	1.32 \pm 0.12	0
1 μ g TBT.g ⁻¹	28.15 \pm 1.59	3.93 \pm 1.88	1.35 \pm 0.21	0
3 μ g TBT.g ⁻¹	27.69 \pm 1.44	3.75 \pm 1.70	1.29 \pm 0.19	0
5 μ g TBT.g ⁻¹	27.34 \pm 1.25	3.60 \pm 1.48	1.24 \pm 0.17	0
8 μ g TBT.g ⁻¹	27.44 \pm 1.80	3.64 \pm 2.13	1.25 \pm 0.24	0
10 μ g TBT.g ⁻¹	26.66 \pm 0.94	3.34 \pm 1.11	1.15 \pm 0.13	0
20 μ g TBT.g ⁻¹	28.89 \pm 1.92	4.22 \pm 2.27	1.45 \pm 0.26	0

Acknowledgements

Mariano Moris (Servicio de Acuario CENPAT) kindly helped with aquaria logistics. This research was partially supported by projects PICT 1232, PICT 2929, PICT 1491, PICT 1476 and PIP 067.

References

- Antizar-Ladislao B (2008) Environmental levels, toxicity and human exposure to tributyltin (TBT)-contaminated marine environment. A review. **Environment International**, 34: 292-308
- Averbuj A, Penchaszadeh PE (2010) On the reproductive biology and impact of imposex in a population of *Buccinanops monilifer* from Mar del Plata, Argentina. **Journal of the Marine Biological Association of the United Kingdom**, 90: 729-734
- Averbuj A, Rocha MN, Zabala S (2014) Embryonic development and reproductive seasonality of *Buccinanops globulosus* (Nassariidae) (Kiener, 1834) in Patagonia, Argentina. **Invertebrate Reproduction and Development**, 58: 138-147
- Axiak V, Vella AJ, Micallef D, Chircop R, Mintoff B (1995) Imposex in *Hexaplex trunculus* (Gastropoda: Muricidae): first results from biomonitoring of tributyltin contamination in the Mediterranean. **Marine Biology**, 121: 685-691
- Bandow C, Weltje L (2012) Development of an embryo toxicity test with the pond snail *Lymnaea stagnalis* using the model substance tributyltin and common solvents. **Science of the Total Environment**, 435-436: 90-95
- Bigatti, G., Cumplido, M. & Averbuj, A. (in press) Gasterópodos de interés comercial en la Provincia del Chubut. Informe para la Mesa Técnica Zona 1 - Subsecretaría de Pesca de

- Chubut - Reglamentación de la pesca de gasterópodos. Inf. N° 30 del LAPEMAR, Laboratorio de Peces y Mariscos de Interés Comercial (CENPAT), 39pp.
- Bigatti G, Primost MA, Cledón M, Averbuj A, Theobald N, Gerwinski W, Arntz W, Morriconi E, Penchaszadeh PE (2009) Biomonitoring of TBT contamination and imposex incidence along 4700 km of Argentinean shoreline (SW Atlantic: From 38S to 54S). **Marine Pollution Bulletin**, 58: 695-701
- Bushong SJ, Jr. LWH, Hall WS, Johnson WE, Herman RL (1988) Acute toxicity of Tributyltin to selected Chesapeake Bay fish and invertebrates. **Water Research**, 22: 1027-1032
- Castellanos Z, Landoni N (1993) Catálogo descriptivo de la malacofauna marina magallánica 9. Neogastropoda Muricidae y Thaisidae. In: Aires CdICdIPdB (ed), La Plata
- Castro LFC, Lima D, Machado A, Melo C, Hiromori Y, Nishikawa J, Nakanishi T, Reis-Henriques MA, Santos MM (2007) Imposex induction is mediated through the Retinoid X Receptor signalling pathway in the neogastropod *Nucella lapillus* **Aquatic Toxicology**, 85: 57-66
- Collin R, Spangler A (2012) Impacts of adelphophagic development on variation in offspring size, duration of development, and temperature-mediated plasticity. **Biological Bulletin**, 223: 268-277
- Commendatore MG, Esteves JL (2007) An assessment of oil pollution in the coastal zone of Patagonia, Argentina. **Environmental Management**, 40: 814-821
- de Mora SJ, Pelletier E (1997) Environmental tributyltin research: past, present, future. **Environmental Technology**, 18: 1169-1177
- Evans SM (1999) Tributyltin Pollution: the Catastrophe that Never Happened. **Marine Pollution Bulletin**, 38: 629-636
- Finney DJ (1971) **Probit analysis: A statistical treatment of the sigmoid response curve**. Cambridge University Press, London
- Foale S (1993) An evaluation of the potential of gastropod imposex as a bioindicator of tributyltin pollution in Port Phillip Bay, Victoria. **Marine Pollution Bulletin**, 26: 546-552
- Garaventa F, Centanni E, Fiorini S, Noventa S, Terlizzi A, Faimali M, Pavoni B (2008) New implications in the use of imposex as a suitable tool for tributyltin contamination: experimental induction in *Hexaplex trunculus* (Gastropoda, Muricidae) with different stressors. **Cell Biology and Toxicology**, 24: 563-571
- Giarratano E, Gil MN, Malanga G (2013) Assessment of antioxidant responses and trace metal accumulation by digestive gland of ribbed mussel *Aulacomya atra atra* from Northern Patagonia. **Ecotoxicology and Environmental Safety**, 92: 39-50
- Giarratano E, Gil MN, Malanga G (2014) Biomarkers of environmental stress in gills of ribbed mussel *Aulacomya atra atra* (Nuevo Gulf, Northern Patagonia). **Ecotoxicology and Environmental Safety**, 107: 111-119
- Gibbs PE, Bryan GW (1986) Reproductive failure in populations of the dog-whelk, *Nucella lapillus*, caused by imposex induced by Tributyltin from antifouling paints. **Journal of the Marine Biological Association of the United Kingdom**, 66: 767-777
- Gibbs PE, Bryan GW (1987) TBT Paints and the demise of the dog-whelk, *Nucella lapillus* (Gastropoda). **Journal of the Marine Biological Association of the United Kingdom**: 1482-1487
- Gil MN, Torres A, Harvey M, Esteves JL (2006) Metales pesados en organismos marinos de la zona costera de la Patagonia argentina continental. **Revista de Biología Marina y Oceanografía**, 41: 167-176
- Gil MN, Torres AI, Commendatore MG, Marinho C, Arias A, Giarratano E, Casas GN (2015) Nutritive and xenobiotic compounds in the alien algae *Undaria pinnatifida* from Argentine Patagonia. **Archives of Environmental Contamination and Toxicology**, 68: 553-565
- González J, Daza E, Vargas C, Cortes C, Guzmán L, Miranda H, Vargas C, Yannicell B (2007) Diagnóstico para la administración y conservación del recurso caracol *Trophon* en Bahía Gente Grande, XII Región. Informe final. **IFOP, Instituto de Fomento Pesquero**, Chile
- Griffin M, Pastorino G (2005) The genus *Trophon* Monfort, 1810 (Gastropoda: Muricidae) in the Tertiary of Patagonia. **Journal of Paleontology**, 79: 296-311
- Horiguchi T, Imai T, Cho HS, Shiraiishi H, Shibata Y, Morita M, Shimizu M (1998) Acute toxicity

- of Organotin compounds to the larvae of the rock shell, *Thais clavigera*, the disk abalone, *Haliotis discus discus* and the giant abalone, *Haliotis madaka*. **Marine Environmental Research**, 46: 469-473
- Horiguchi T, Nishikawa T, Ohta Y, Shiraishi H, Morita M (2007) Retinoid X receptor gene expression and protein content in tissues of the rock shell *Thais clavigera*. **Aquatic Toxicology**, 84: 379-388
- Horiguchi T, Ohta Y, Nishikawa T, Shiraishi F, Shiraishi H, Morita M (2008) Exposure to 9-retinoic acid induces penis and *vas deferens* development in the female rock shell, *Thais clavigera*. **Cell Biology and Toxicology**, 24: 553-562
- IMO (2002) International Maritime Organization. Focus on IMO-Anti-fouling systems. International Convention on the Control of Harmful Anti-fouling Systems on Ships. Resolution A.928(22). <http://www.imo.org>
- Laughlin RB, Jr., Thain JE, Davidson BM, Valkirs AO, Newton FC, III (1996) Experimental studies of chronic toxicity of Tributyltin compounds. In: Champ MA, Seligman PF (eds) **Organotin**. Springer Netherlands, p 191-217
- Lee J, Lee K-T, Park GS (2005) Acute toxicity of heavy metals, Tributyltin, ammonia and polycyclic aromatic hydrocarbons to benthic amphipod *Grandidierella japonica*. **Ocean Science Journal**, 40: 61-66
- Leiva GE, Castilla JC (2002) A review of the world marine gastropod fishery: evolution of catches, management and the Chilean experience. **Reviews in Fish Biology and Fisheries**, 11: 283-300
- Matthiessen P (2013) The prospects for routine testing of chemicals for endocrine-disrupting properties and potential ecological impacts **Endocrine Disruptors**. John Wiley & Sons, Inc., p 373-384
- McClellan-Green P, Romano J, Rittschof D (2006) Imposex induction in the mud snail, *Ilyanassa obsoleta* by three Tin compounds. **Bulletin of Environmental Contamination and Toxicology**, 76: 581-588
- Meador JP (2000) Predicting the fate and effects of tributyltin in marine systems. **Reviews of Environmental Contamination and Toxicology**, 166: 1-48
- Moore DW, Dillon TM, Suedel BC (1991) Chronic toxicity of tributyltin to the marine polychaete worm, *Neanthes arenaceodentata*. **Aquatic Toxicology**, 21: 181-198
- Morris HR (1987) Acute toxicity of Tributyltin chloride to embryos and larvae of two bivalve mollusks, *Crassostrea virginica* and *Mercenaria mercenaria*. **Bulletin of Environmental Contamination and Toxicology**, 39: 1012-1019
- Narvarte MA (2006) Biology and fishery of the whelk *Buccinanops globulosum* (Kiener, 1834) in northern coastal waters of the San Matías Gulf (Patagonia, Argentina). **Fisheries Research**, 77: 131-137
- Oberdörster E, McClellan-Green P (2000) The neuropeptide APGWamide induces imposex in the mud snail, *Ilyanassa obsoleta*. **Peptides**, 21: 1323-1330
- Oberdörster E, McClellan-Green P (2002) Mechanisms of imposex induction in the mud snail, *Ilyanassa obsoleta*: TBT as a neurotoxin and aromatase inhibitor. **Marine Environmental Research**, 54: 715-718
- Oberdörster E, Romano J, McClellan-Green P (2005) The neuropeptide APGWamide as a penis morphogenic factor (PMF) in gastropod mollusks. **Integrative and Comparative Biology**, 45: 28-32
- Oehlmann J, Fioroni P, Stroben E, Markert B (1996) Tributyltin (TBT) effects on *Ocenebrina aciculata* (gastropoda: Muricidae): Imposex development, sterilization, sex change and population decline. **Science of the Total Environment**, 188: 205-223
- Oehlmann J, Schulte-Oehlmann U, Tillmann M, Markert B (2000) Effects of endocrine disruptors on prosobranch snails (Mollusca: Gastropoda) in the laboratory. Part I: Bisphenol a and octylphenol as xeno-estrogens. **Ecotoxicology**, 9: 383-397
- Panagoula B, Panayioti M, Iliopoulou-Georgoudaki J (2002) Acute toxicity of TBT and Irgarol in *Artemia salina*. **International Journal of Toxicology**, 21: 231-233
- Pastorino G (1993) The taxonomic status of *Buccinanops d'Orbigny*, 1841 (Gastropoda: Nassariidae). **The Veliger**, 36: 160-165
- Pastorino G (2005) A revision of the genus *Trophon* Montfort, 1810 (Gastropoda: Muricidae) from southern South America. **The Nautilus**, 119: 55-82
- Penchaszadeh PE, Averbuj A, Cledón M (2001) Imposex in gastropods from Argentina (South-Western Atlantic). **Marine Pollution**

- Bulletin**, 42: 790-791
- Pessoa MF, Fernando A, Oliveira JS (2001) Use of imposex (pseudohermaphroditism) as indicator of the occurrence of organotin compounds in Portuguese coastal waters - Sado and Mira estuaries. **Environmental Toxicology**, 16: 234-241
- Ruiz JM, Bryan GW, Gibbs PE (1994) Chronic toxicity of water tributyltin (TBT) and copper to spat of the bivalve *Scrobicularia plana*: ecological implications. **Marine Ecology Progress Series**, 113: 105-117
- Ruiz JM, Bryan GW, Gibbs PE (1995) Acute and chronic toxicity of tributyltin (TBT) to the pediveliger larvae of the bivalve *Scrobicularia plana*. **Marine Biology**, 124: 119-126
- Santos MM, Castro LFC, Vieira MN, Micael J, Morabito R, Massanisso P, Reis-Henriques MA (2005) New insights into the mechanism of imposex induction in the dogwhelk *Nucella lapillus*. **Comparative Biochemistry and Physiology, Part C**, 141: 101-109
- Spooner N, Gibbs PE, Bryan GW, Goad LJ (1991) The effect of tributyltin upon steroid titres in the female dogwhelk, *Nucella lapillus*, and the development of imposex. **Marine Environmental Research**, 32: 37-49
- Thain J, Waldock M, Waite M (1987) Toxicity and degradation studies of Tributyltin (TBT) and Dibutyltin (DBT) in the aquatic environment **OCEANS '87 1398-1404**,
- Waldock MJ, Thain JE, Waite ME (1987) The distribution and potential toxic effects of TBT in UK estuaries during 1986. **Applied Organometallic Chemistry**: 287-301
- Zhang K, Shi J, He B, Xu W, Li X, Jiang G (2013) Organotin compounds in surface sediments from selected fishing ports along the Chinese coast. **Chinese Science Bulletin**, 58: 231-237

Received: April 2015


Accepted: June 2015

Published on-line: August 2015



ANNEX I from: Acute toxicity in two edible marine gastropods with different sensitivity to Tributyltin

M.A. PRIMOST *ET AL.*



ANNEX 1 Figure 1.- Body mass weight vs total weight and size vs total weight for *Trophon geversianus* and *Buccinanops globulosus* from Cerro Avanzado".